

IODOFLUOROALKANES: PROPERTIES, SYNTHESIS, APPLICATION

A. V. Morozov, B. N. Maximov

Federal State Unitary Enterprise "Russian Scientific Centre (RSC) "Applied Chemistry"

The increased interest for iodofluoroalkanes has been created because of their unique properties, which define their value in terms of practice. Since 1967 the studies regarding synthesis of iodofluoroalkanes had been being conducted in RSC of Applied Chemistry. In the present review we have presented the main properties and fields of application for iodofluoroalkanes, the results of studies regarding the development of methods and obtaining technology for iodofluoroalkanes conducted by the authors.

1. Some Physical and Chemical Properties of Iodofluoroalkanes

Iodofluoroalkanes - heavy colourless compounds with specific odour and weak narcotic properties. Lowest members of homologous series - gases and liquids, homologues with the number of carbon atoms over 8 - waxes, with the number of atoms over 40 - outwardly remind polytetrafluoroethylene [1].

The density of iodofluoroalkanes is higher than the density of corresponding iodoalkanes, which is because of greater mass of fluorine atoms compare to mass of hydrogen atoms in molecules; it varies within the limits of 2- 2,7 g/cm³. The boiling points of iodofluoroalkanes are on average 60°C lower than the ones of corresponding iodoalkanes through smaller intermolecular interaction caused by the absence of hydrogen bonds typical for iodoalkanes.

Iodofluoroalkanes as well as fluorocarbons are characterized by low surface tension, high vapour pressure, small evaporation energies. Iodofluoroalkanes are characterized by strong continuous absorption in short-range UV field. The high value of ionization potentials compare to iodoalkanes (1,25 - 1,4 eV) shows higher delocalization of iodine atoms free electrons through the whole [2]. In consequence of electron-seeking influence of fluorine atoms in the molecule of iodofluoroalkane on the carbon atom connected to iodine a perceptible positive charge is formed, which is partly transferred on easy to polarize iodine atom, and as a result the electron density in the zone of C-I bond is decreasing as well as the value of bond opening energy [3].

Iodofluoroalkanes are not flammable in contrast to iodoalkanes; they dissolve in organic solvents better compare to fluorocarbons. They practically do not dissolve in water. They are moderate toxic compounds. α,ω - Diiodoperfluoroalkanes remind monoiodofluoroalkanes in terms of their properties, however their density and boiling points are significantly higher than the ones of corresponding monoiodides.

Lowest members of diiodides series - heavy colourless liquids, the highest ones - waxy compounds [1]. C-I bonds of lowest α,ω - diiodoperfluoroalkanes are less durable than the ones of monoiodofluoroalkanes, but as the CF₂I groups are moving away from each other the C-I bond in terms of durability becomes the same as the one of monoiodofluoroalkanes [4].

Both mono- and diiodoperfluoroalkanes turn rose at the light exposition as a result of photo-chemical decomposition with iodide isolation. In terms of their chemical properties iodofluoroalkanes significantly differ from iodoalkanes because of the combination of high inert fluoroalkyl group and reactive iodide in the molecule. It is considered, that C-I bond in iodofluoroalkanes loses its ability for heterolytic decomposition [5]. Because of the partial positive charge in the iodide atom iodofluoroalkanes are not inclined to nucleophilic

iodide replacement - the most important type of reaction for [6]; however the nucleophilic substitution is observed at thermal initiation or UV irradiation during reaction with alcohols, ammonia, water with forming of hydrofluoroalkanes [7].

Iodo fluoroalkanes are resistant to hydrolysis, but during the boiling together with the alkali aqueous solutions produce hydrofluoroalkanes [8]. Iodine is being substituted by fluorine in iodo fluoroalkanes at influence of strong fluorinating agents - F_2 , CoF_3 , BrF_3 , SbF_5 [9], medium strong fluorinating agents (JF_5 , SbF_3Cl_2) are able to replace iodide for fluorine only in the lowest α,ω - diiodo fluoroalkanes with forming of monoiodo fluoroalkanes [5].

The substitution of iodine for Cl or Br is possible only under the conditions of free-radical reaction [10]. Iodo fluoroalkanes are oxidation resistant at routine temperatures, but under the harsh conditions they are oxidized to carboxylic acids under the influence of concentrated H_2SO_4 , HNO_3 , H_2O_2 , $KMnO_4$ [11].

2. Iodo fluoroalkanes' Application Fields

The majority of iodo fluoroalkanes' known reactions are free-radical, in which the C-I bond is subject to homolysis during heating, radiation or attack of free radicals. The main field of application for the lowest iodo fluoroalkanes - as telogens at telomerization with forming of products used for the synthesis of fluoroacids [11], fluoroalcohols [12], monomers for thermo- and chemically stable polymers [12], at modification of polymers, obtaining of lubricating oils, dielectrics [10] etc.

Fluoroalkylation of aliphatic, aromatic, heterocyclic, elementorganic compounds allowing obtaining the compounds of new precious characteristics [13- 15] is an intensively developing field of application for iodo fluoroalkanes.

The use of fluoroalkanes in laser systems (optical laser oscillators) is a new direction, which is caused by the ability of iodo fluoroalkanes to photodissociation according to C-I bond with forming of high concentration of iodide excited atoms much higher than iodoalkanes can produce [16]. Taking into account the publications on lasers using iodo fluoroalkanes and of high power and coefficient of efficiency, this field of their application is becoming important and holds much promise [17].

Lowest monoiodo fluoroalkanes find their limited application as highly effective extinguishers for the volumetric fire extinguishing, computer techniques, different types of documentation due to their high thermal stability, high density of vapours, and low toxicity [18]. Due to their insignificant ozone hazard some of iodo fluoroalkanes (C_2F_5I etc.) are used as refrigerants in small refrigerating units and as excellent solvents for cleaning the goods of electronics and precision technics [18, 19].

3. Main Obtaining Methods For Fluoroalkanes; Advantages and Disadvantages

Historically the main obtaining method of iodo fluoroalkanes was the replacement of carboxylic group in acids with halogen (iodide), though later iodo fluoroalkanes were obtained by the reaction of Hunsdiecker at decarboxylation of fluorocarboxylic acid silver salt with iodide [20, 21]. The chance to obtain iodo fluoroalkanes of $CF_3(CF_2)_nI$ row should be referred to advantages of the method, though the high cost of initial materials, low productivity due to the long period of the process, technological inefficiency, the significant quantity of by-products (10-26% have transferred the method to preparatory class.

The method of sodium and potassium salts decarboxylation in the medium of DMF at 150 - 160°C in autoclave [22] is to be known, however the duration of periodical process and significant quantity of by-products (20-30%) bound the introduction of this method.

Taking into account the availability and comparatively low cost of the initial raw materials the methods oriented to fluoroolefines appeared to be mostly progressive. The methods of iodofluorination of fluoroolefines using the mixture of iodine and JF_5 over catalysts got most famous, at that the ratio of J_2 and JF_5 was regulated as 2 : 1, which corresponded to the postulated forming at heating up the reagents of JF monofluoride, which produced iodofluoroalkane adding by double bond of olefine [24 - 26].

(Ag, Cu, Sn, Sb, Cd, Co, Pb, Ni, Cr) metals [25] were used as catalysts. According to the methods mentioned above the synthesis process was carried out at self developing pressure in autoclaves and at high temperatures (up to 150-225°C). The small productivity caused by the limits of reagents' feeding, long period of the process, instability of the process parameters due to both its periodicity and difficulties of abstraction the heat of exothermal reactions, particularly the telomerization reactions following high heat emission were the methods' disadvantages.

The continuous synthesis method of $\text{C}_2\text{F}_5\text{J}$ directly from J_2 , F_2 and C_2F_4 , is described, in it the gaseous mixture of fluorine and iodine reacts at 150°C forming JF_5 , which flows down to the column with nozzle, reacting with the mixture of iodine and C_2F_4 vapours coming from below; the method has not spread because of low yields as well as its potential explosibility [28].

A certain point of view regarding the mechanism of fluoroolefines' iodofluorinating using mixture of iodine and JF_5 was not to find in the scientific literature. The forming of JF out of the mixture of iodine and JF_5 during the process of iodofluorination postulated by the number of authors did not conform the information of Ruffo and Braida on the absence of JF in such system at room temperature and higher temperatures [29].

Analogously, Schmeisser, Scharf and Sartori [30, 31] stated, that JF as a substance was stable only at low temperatures (under -14°C) and at heating above -14°C dissociated for J_2 and JF_5 . Taking into account the high temperature of iodofluorination of the mentioned methods [24 - 27], which exceeds the JF temperature of stability, its existence in the processes is unlike to be true. Taking into account the conducted analysis of iodofluoroalkanes synthesis methods described in literature, we developed a method and technology for obtaining of iodofluoroalkanes free from the minuses of known methods, at that the major attention was drawn to iodofluoroalkanes perspective for laser systems using.

4. The Study Of α,ω - Diiodofluoroalkanes Synthesis Process

First of all we have studied the thermally initiated reaction of iodine and C_2F_4 for the purpose of obtaining the 1,2- diiodtetrafluoroethane, out of which it was possible to obtain quantitatively (as we stated) $\text{C}_2\text{F}_5\text{J}$, which has got a potential for application in laser systems. Although there are some reactions of iodine and C_2F_4 initiated photochemically, catalytically or by heating [32 - 34] described in literature, the processes have been stated scarcely, they are characterized by instability and low productivity. We have proved, that gas-cycle reaction of iodine and C_2F_4 in stationary system passes significantly only above 110°C.

Its dependence on inhibitors, on the size of specific area of the reactor, on temperature, which initiated homolysis of molecular iodine into atoms, eased by small amount of energy of J-J bond - 150,7 KJ/mole pointed on radical character of the reaction. The jump in the rate of reaction at increasing the temperature conformed the increasing disintegration of the iodine molecules into atoms which goes with exponential rate constant [16]. S-formed kinetic C_2F_4 consumption curve at the reaction with iodine looked like the curves of autocatalytic reactions, which was due not to the branching of kinetic chain at the dissociation of formed diiodides into radicals, but to the appearance of

the liquid phase out of 1,2-diiodofluoroalkane, during which the reaction was passing at a higher rate.

The studies of iodine and C_2F_4 reaction in the medium of low-polarity solvents, including $C_2F_4J_2$, allowed proving, that the reaction as a radical chain process was going at as low as 60-100°C; as we can ignore the thermal homolysis of iodine at that temperatures (the equilibrium constant of the $J_2 \rightarrow 2J$ reaction is equal to $1,33 \cdot 10^{-16}$ at 100°C), the initiation of radical process, as we had thought was going according to the mechanism of molecularly induced homolysis at donor-acceptor interaction of diluted molecules of iodine and C_2F_4 .

The shift of absorption band of the double bond of C_2F_4 in IR-spectrum (till 1665 cm^{-1}) at introduction of iodine into C_2F_4 solution tells us about that as well as the appearance of the stretching vibrations band of polarized molecule of J_2 at 206 cm^{-1} . Weakened at that interaction J-J bond broke at lower temperature within the range of 60-100°C forming radicals, initiating forming of α,ω -diiodofluoroalkanes.

According to the data obtained by our specialists during kinetic studies [36], the rate of interaction of iodine and C_2F_4 can be described by the equation:

$$r = K_{\text{effective}} \cdot [C_2F_4]^{3/2} \cdot [J_2]^{1/2}$$

where $K_{\text{effective}}$ is a function of constants of the rates of elementary reactions of initiation, development and break of chains;

The discovered regularities allowed developing the obtaining method for 1,2-diiodotetrafluoroethane [35], and also the obtaining method for α,ω -diodoperfluoroalkanes [36].

5. The Study of Monoiodofluoroalkanes Synthesis

Iodofluorination using the mixture of J_2 - JF_5 fluorolefines was studied by our specialists as the most promising direction in the development of iodofluoroalkanes obtaining method. We have chosen the representative of "hard" Lewis acids - SbF_3 among accelerating metals fluorides iodo-fluorination process, it provides high rate for the process, as well as it is available and convenient in use. We have proved, that under certain conditions SbF_3 in the solution of J_2 in JF_5 formed complex anions of distorted octahedral symmetry connected by bridge fluorine- $Sb_nF_{5n+1}^-$, at that the transfer of SbF_3 from typical for it p^3 condition into sp^3d^2 hybrid condition. At acceptor influence of these anions on the easily polarized J_2 molecule the molecular cation of J_2^+ has formed, it enters the composition of the complex of $J_2^+Sb_nF_{5n+1}^-$, mainly $J_2^+Sb_2F_{11}^-$ (according to the elemental analysis).

The analysis of IR- and NMR- spectra of such formations confirmed the structure of the complex [36, 37]: the presence of band at 240 cm^{-1} in IR-spectrum from molecular cation J_2^+ , bands at 578 and 692 cm^{-1} from anions $Sb_2F_{11}^-$, bands from bridge bond's vibrations of Sb-F-Sb at 510 cm^{-1} and ($J_2^+ \dots F-Sb^-$) at 490 cm^{-1}

bands of degenerate vibrations from four fluorine ligands at each stibium atom;

Sb-F-Sb - $\nu_{\text{as}}(SbF_4)$ at 665 cm^{-1} and $\nu_s(SbF_4)$ at 530 cm^{-1} and 545 cm^{-1} ; bands at 620 cm^{-1} from fluorine in *trans* position relatively F-Sb-F bridge.


The generating of J_2^+ cations in J_2 - JF_5 iodofluorinating system - the antimony fluorides

created the conditions for conducting the synthesis of iodo fluoroalkanes at iodo fluorinating of fluoroolefines in flow system at moderate temperatures and pressure close to the atmospheric one.

At disappearance of J^{2+} cations in the J_2 - JF_5 - SbF_3 system for example at anhydrous hydrogen fluoride's getting into it, the rate of iodo fluorination of fluoroolefine sharply decreased. The study of the J_2 - JF_5 - SbF_3 -HF system using methods of IR- and NMR-spectroscopy revealed the solvation of antimony anions by HF molecules, which hampered their acceptor influence on iodine and formation of J^{2+} cations.

It confirmed the disappearance of the band at 240 cm^{-1} in IR-spectrum at introduction of HF into iodo fluorinating system, the disappearance of the blue tone of solutions, which appeared at the presence of iodine cations in solutions, the alterations of chemical shifts in NMR- F^{19} spectrum and of signals forms of F^{19} in the $Sb_2F_{11}^-$ anion [36, 37].

Solvation of $Sb_2F_{11}^-$ anions using HF molecules and lowering of catalytic activity of antimony fluorides is reversible: at elimination of HF out of reaction mixture the catalyst restores its activity. The stated connection between the iodo fluorinating rate and the rate of fluoroolefines' with J^{2+} cations transformation points out their participation in iodo fluorination, and the decreasing of fluoroolefines reactivity at iodo fluorination in the J_2 - JF_5 - SbF_3 system, according to the decreasing the electron density of double bonds was the sign of the original electrophilic addition of J^{2+} cation to fluoroolefine - the fluorinating rate decreased in the row:




(as the reactivity towards the electrophiles)

The constant of C_2F_4 iodo fluorination rate defined by us significantly exceeded the one during the C_3F_6 iodo fluorination, and the energy of activation was by 8 kJ/mole lower [36].

SbF_5 and mixed antimony chorofluorides act at iodo fluorination like SbF_3 , though smaller selectivity and rates of iodo fluorination were caused by HF admixture to that associate fluorides, the purification is rather difficult [36].

The interaction of J^{2+} and fluoroolefine begins probably with the formation of π -complex during the equilibrium process, that role of that complex is in the spatial orientation of reagents into the position necessary for reaction. Then the electrophilic atom of iodine cation J^{2+} forms together with olefin π - electron pair a carbocation or a 3-member iodonium cation. After the addition of F^- anion-fluoride and neutralization of charge the iodo fluoroalkane forms.



The fact, that isomers form during iodo fluorination of fluoroolefines with different substitutes at the double bond ($CF_2=CFCl$ give the mixture of 45% $CF_3-CFClI$ and 55% CF_2Cl-CF_2I) is the argument for iodonium cation as an intermediate formation.

As one side in the iodonium cation is shielded with the iodine bridge, nucleophile F^- can attach to any of the two atoms of iodonium cycle carbon from the opposite side. In C_3F_6

the electrophilic atom of iodine is shifting to the carbon of higher electron density and the orientation of F^- anion addition is definite: CF_3-CF_2-I is formed. The role of JF_5 is in the completing of fluorine ligands of Sb atoms being spent. The synthesis of iodo fluoroalkanes in flow system at sparging of fluorolefines through the iodine and SbF_3 solution in JF_5 was going at creation of saturated solutions of iodine and antimony fluorides; due to the increased rate of reaction the mass exchange processes began to influence the synthesis and could limit the rate of iodo fluoroalkanes forming.

As the mass transfer of fluoroolefine out of gas bubbles into solution was the limiting stage of synthesis according to our statement [38, 40], the creating of developed interphase surface was necessary for the transferring of synthesis process into kinetic field. It should be noticed, that the synthesis of iodo fluoroalkanes under the conditions of barbotage virtually had not been accompanied by the formation of telomers due to the outing of formed products from the reaction zone and low, close to the atmospheric pressure during synthesis.

At that, the evaporating heat isolation system provided keeping of isothermal conditions of the synthesis process; taking into account all the heat expenses, the heat effect had not exceeded 0,8 - 7 kJ/mole, while under the static reaction conditions the heating up and decomposition of the mixtures due to the difficulty of heat outing/isolation had been observed [27]. The studies regarding the updating of iodo fluoroalkanes synthesis technology in the flow system allowed developing of the mathematical model of C_2F_4 and C_3F_6 iodo fluorinating processes; the solubility of reagents in JF_5 , mass-exchange coefficients, the dependence of interphase surface on the barbotage regime were defined experimentally. Under the simplifying conditions of quasistationary conditions the kinetics and mass-exchange were described.

As a result the mathematical model of pentafluoroiodoethane and 2-iodoheptafluoropropane [40] synthesis was composed; under the operational conditions of pilot plant its testing (the model) proved, that the model adequately described the synthesis process of iodo fluoroalkanes synthesis.

The results obtained during the study of fluorolefines iodo fluorination allowed developing of the iodo fluoroalkanes obtaining method, which was more up-to-date, than the known ones [39] and as a result the obtaining technologies of pentafluoroiodoethane, 2-iodoheptafluoropropane, trifluoromethoxytetrafluoroiodoethane were developed and introduced at Pilot Plant of RSC "Applied Chemistry". Below you will find the characteristics of obtained iodo fluoroalkanes (Table 1).

Table 1. Physical Characteristics of Synthesized Iodo fluoroalkanes

Compounds	Boiling point (mm Hg), °C	Melting point (mm Hg), °C	d_4^{20} kg/m ³
CF_3-CF_2-I	12,5 (760)	-117	2080
CF_3CFH-I	39,4 (760)	-98	2070
$CF_3CFCI-I$	56 (760)	-76	2152
CF_2Cl-CF_2I	56,4 (760)	-75,3	2157
CF_3CH_2I	55 (760)	-87	1992
$(CF_3)_2CFI$	40 (760)	-65	2063

CF ₃ -CF ₂ CF ₂ CF ₂ J	66 (760)	-72	2012
CF ₃ CF ₂ CFJ-CF ₃	65,5 (760)	-57	2020
CF ₃ -(CF ₂) ₅ J	117 (760) 84 (290)		2012
CF ₃ -(CF ₂) ₇ J	160 (760)		2004
CF ₃ -(CF ₂) ₉ J	195 (760)	66	1998
CF ₃ -(CF ₂) ₁₁ J	110 (18)		1997
CF ₃ OCF ₂ CF ₂ J	46,5 (760)	-79	2015
C ₃ F ₇ OCF ₂ CF ₂ J	86,5 (760)		1963
(CF ₃) ₂ CJ-CFO	87 (760)	51	2024
(CF ₃) ₂ CF-CF ₂ CF ₂ J	94 (760)	-50	2031
(CF ₃) ₂ CF-(C ₂ F ₄) ₂ J	138 (760)	-38	2028
(CF ₃) ₂ CF(C ₂ F ₄) ₃ J	174 (760)		2008
J-CF ₂ -CF ₂ -J	112,5 (760)	-43	2630
J-(CF ₂ CF ₂) ₂ -J	150 (760)	-9	2487
J(CF ₂ -CF ₂) ₃ -J	185 (760)	25	2357
J(CF ₂ CF ₂) ₄ -J	216 (670)	76	2259
J-CF ₂ -CF ₂ Br	88 (760)	-51	2380

References

Journal contents